

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/62899 A2**

(51) International Patent Classification<sup>7</sup>: C12N 5/06, 5/00

(21) International Application Number: PCT/US01/05252

(22) International Filing Date: 20 February 2001 (20.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/510,444 21 February 2000 (21.02.2000) US

(71) Applicant: WISCONSIN ALUMNI RESEARCH  
FOUNDATION [US/US]; 614 Walnut Street, P O Box  
7365, Madison, WI 53707-7365 (US).

(72) Inventors: THOMSON, James, A.; 1807 Regent Street,  
Madison, WI 53705 (US). MARSHALL, Vivienne,  
S.; 4326 Buckeye Road, Madison, WI 53716 (US).  
SWIERGIEL, Jennifer, J.; 10219 Hawks Point Trail,  
Roscoe, IL 61073 (US).

(74) Agent: SEAY, Nicholas, J.; Quarles & Brady LLP, P O  
Box 2113, Madison, WI 53701-2113 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: METHOD OF MAKING EMBRYOID BODIES FROM PRIMATE EMBRYONIC STEM CELLS

(57) Abstract: Primate embryoid bodies are formed from primate ES cells. The ES cells form clumps. One then removes the clumps, as clumps, and permits incubation under non-adherent conditions. The development of embryoid bodies from primate ES cells is dependent on maintaining the aggregation of cells, as individualized cells will rapidly die.

WO 01/62899 A2

METHOD OF MAKING EMBRYOID BODIES FROM  
PRIMATE EMBRYONIC STEM CELLS  
CROSS REFERENCES TO RELATED APPLICATIONS

Not applicable.

5 STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

The work with the non-human primate cells described below was supported by a grant with United States government support awarded by the following agency: NIH RR11571. The United States has certain rights in this invention. No U.S. government funds were used for the work with human cells described herein.

BACKGROUND OF THE INVENTION

Undifferentiated primate embryonic stem ("ES") cells can be cultured indefinitely and yet maintain the potential to form differentiated cells of the body. See 15 U.S. patent 5,843,780; J. Thomson, et al., 282 Science 1145-1147 (1998); and J. Thomson, et al., 38 Biology 133-165 (1998). The disclosure of these publications and of all other publications referred to herein are 20 incorporated by reference as if fully set forth herein.

Primate ES cells thus provide an exciting new model for understanding the differentiation and function of human tissue, and offer new strategies for drug discovery and testing. They also promise new therapies based on 25 the transplantation of ES cell-derived tissues. For example, human and rhesus monkey ES cells injected into immunocompromised mice form benign teratomas with advanced differentiated derivatives representing all three embryonic germ layers. Easily identified 30 differentiated cells in human ES cell teratomas include smooth muscle, striated muscle, bone, cartilage, gut and respiratory epithelium, keratinizing squamous epithelium, hair, neural epithelium, and ganglia.

Human and non-human primate ES cell lines provide a particularly powerful new model for understanding normal human development and thus also for understanding 35

abnormal human development. Because of the potential risk to the resulting child, experimental manipulation of the post-implantation human embryo is ethically unacceptable and as a result functional studies on human embryos are lacking. Consequently, what is known about human development in the early post-implantation period is based almost entirely on static histological sections of a few human embryos and on analogy to experimental embryology studies of the mouse.

However, early mouse and primate development differ significantly. For example, human and mouse embryos differ in the timing of embryonic genome expression, in the formation, structure, and function of the fetal membranes and placenta and in the formation of an embryonic disc instead of an egg cylinder. The earliest events of human development are critically involved in human infertility, pregnancy loss, and birth defects. Primate ES cells offer a new window for understanding these early human developmental events and for understanding the pathogenesis of developmental failures.

Primate ES cells also provide a potentially unlimited source of differentiated, euploid, non-transformed cells for investigators interested in the normal function and pathology of specific differentiated primate cells. Such purified populations of specific ES cell-derived cells will also likely be useful for drug discovery, toxicity screens, and will provide a source of cells for transplantation.

For tissues such as the heart that completely lack a tissue-specific stem cell, primate ES cells will prove even more valuable. Primate ES cells also offer the promise of new transplantation therapies. When disease results from the destruction or dysfunction of a limited number of cell types, such as in Parkinson's disease (dopaminergic neurons), or juvenile onset diabetes

mellitus (pancreatic  $\beta$ -islet cells), the replacement of those specific cell types by ES cell derivatives could offer potentially life long treatment.

To accomplish these goals, it is desirable to more efficiently differentiate ES cells to specific lineages. Considerable progress in causing non-primate ES cell differentiation to neural, hematopoietic, and cardiac tissue has been made. See e.g. T. Doetschman, et al., 87 J. Embry. And Exper. Morph. 27-45 (1985); G. Keller, 7 Current Op. In Cell Biol. 862-869 (1995); U.S. patent 5,914,268. In each of these examples, ES cells were first formed into "embryoid bodies", three-dimensional ES cell aggregates that facilitate subsequent differentiation.

However, analogous experiments on primate ES cells demonstrated that embryoid body formation by conventional murine protocols fail. In such conventional protocols ES cells are dispersed to single cells, and either allowed to aggregate into embryoid bodies under conditions that prevent cell attachment to the substrate, or the ES cells are allowed to grow into embryoid bodies from single cells or clusters suspended in methylcellulose. We have learned that primate ES cells die rapidly when dispersed to single cells if attachment is prevented, so they do not successfully aggregate, and they therefore do not grow out from clones in methylcellulose.

It can therefore be seen that a need exists for improved methods for producing primate embryoid bodies, and differentiated cells derived therefrom.

#### BRIEF SUMMARY OF THE INVENTION

The present invention provides a method for producing primate embryoid bodies from colonies of primate embryonic stem cells that are adhering to a substrate. One removes the adhering colonies of the embryonic stem cells from the substrate in clumps. One

then incubates the clumps in a container under conditions in which the clumps are essentially inhibited from attaching to the container and coalesce into embryoid bodies. For purposes of this application, a clump is a  
5 grouping of two or more stem cells, preferably a clump large enough to be visible to the naked eye.

In one preferred form the removal step is in the presence of an agent that promotes disassociation of the clumps from the substrate as clumps. A purely chemical  
10 agent such as Versene® calcium disodium EDTA chelating agent can be used. However, more preferred is a proteinase that preferentially acts on the extra cellular matrix such as dispase, collagenase, catalase, neuraminidase, pancreatin, pancreatic elastase or  
15 trypsin. If trypsin is used the removal step must be conducted quickly and at relatively low concentrations in order to prevent the trypsin from also destroying the clumps. Enzyme EDTA mixes can also be used to advantage.

In another form the removal step involves  
20 mechanically scraping the clumps from the substrate as clumps.

In another aspect the incubation step can be conducted by agitating the container (e.g. by gently rocking, shaking, or vibrating it), the container for the  
25 incubation step can be a non-attaching bacterial grade culture plastic, and/or the incubation step can be in the presence of a serum-free medium which lacks serum attachment factors.

In another aspect the invention provides primate  
30 embryoid bodies that have been derived (directly or indirectly) using the above methods.

In still another aspect the invention provides differentiated cells derived (directly or indirectly) from the embryoid bodies.

35 In accordance with the present invention, primate ES

cells that have been cultured under standard conditions (see e.g. U.S. patent 5,843,780) are permitted to overgrow, pile up and/or otherwise closely associate in clumps on a substrate (e.g. a plastic tissue culture plate with standard feeder layer). They are then removed as clumps from the substrate (e.g. by incubating the colonies with an enzyme which attacks the ES cell colony's attachment to the substrate more strongly than ES cell attachments with ES cells). In such a case the enzyme could be dispase at a concentration of about 10 mg/ml.

Alternatively, the clumps could be removed as clumps by mechanically scraping with a micropipette, cell scraper, or the like.

The essentially intact colonies are then incubated under non-attaching conditions (preferably continuous rocking of the culture dish, culture on non-attaching bacterial grade culture plastic, and/or continuous culture in the presence of serum-free medium which lacks serum attachment factors). The colonies can then quickly coalesce into compact embryoid bodies, which can thereafter be allowed to differentiate either in continuous suspension, or after re-attachment to a substrate. Such embryoid bodies can be used to derive differentiated derivatives of endoderm, mesoderm, and ectoderm, and for obtaining other desired lineages.

It is an advantage of the present invention that it provides effective methods of forming primate embryoid bodies from primate embryonic stem cell lines. Another advantage of the present invention is to provide primate embryoid bodies suitable for differentiation into other primate cell types. Other features and advantages of the present invention will become apparent after study of the specification and claims which follow.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Formation Of Embryoid Bodies

Primate embryonic stem cells (e.g. rhesus or human -  
U.S. patent 5,843,780; J. Thomson, et al., 282 Science  
1145-1147 (1998)) are cultured on mitotically inactivated  
(3000 rads g-radiation) mouse embryonic fibroblasts,  
prepared at  $5 \times 10^4$  cells/cm<sup>2</sup> on tissue culture plastic  
previously treated by overnight incubation with 0.1%  
gelatin. E. Robertson, Embryo-derived Stem Cell Lines.  
In: Teratocarcinomas And Embryonic Stem Cells: A  
Practical Approach IRL Press: Washington, D.C., 71-112  
(1987). Culture medium consists of 79% Dulbecco's  
modified Eagle medium (DMEM; 4500 mg of glucose per  
liter; without sodium pyruvate), 20% fetal bovine serum  
(FBS), 0.1 mM 2-mercaptoethanol, 1 mM L-glutamine and 1%  
nonessential amino acid stock (GIBCO).

One allows colonies to form clumps over a period of  
hours. ES cell colonies can then be removed from the  
tissue culture plate using physical or chemical methods  
that keep the ES cells in clumps.

For dispase or collagenase removal of ES cell  
colonies from the culture plate, the culture medium is  
removed from the ES cells. Dispase (10 mg/ml in ES  
culture medium) or collagenase (1 mg/ml solution in DMEM  
or other basal medium) is added to the culture plate. The  
culture plates are returned to the incubator for 10-15  
minutes.

After dispase treatment the colonies can either be  
washed off the culture dishes or will become free of the  
tissue culture plate with gentle agitation. After  
collagenase treatment the cells can be scraped off the  
culture dish with a 5 ml glass pipette. Some dissociation  
of the colonies occurs, but this is not sufficient to  
individualize the cells. After chemical removal of the  
cells from the tissue culture plate, the cell suspension

is centrifuged gently for 5 minutes, the supernatant is removed and discarded, the cells are rinsed, and the cells are resuspended in culture medium with or without serum.

5           Mechanical removal of the cells is achieved by using a pulled glass pipette to scrape the cells from the culture plate. Cell clumps can be immediately resuspended, without centrifugation, in fresh tissue culture medium.

10           Once colonies are removed from the tissue culture plate, the ES cells should remain in suspension during further embryoid body formation. This can be achieved by, for example, gently and continuously rocking the cell suspension. Cell suspensions are aliquoted into wells of  
15   6-well tissue culture dishes, placed inside a sealed, humidified isolation chamber, gassed with 5% CO<sub>2</sub>, 5% O<sub>2</sub> and 90% N<sub>2</sub> and placed on a rocker (Red Rocker, Hoefer Scientific Instruments). The rocker is housed inside an incubator maintained at 37°C. The culture plates can be  
20   rocked continuously for at least 48 hours and up to 14 days.

          Every 2 days the plates are removed from the rocking device, the culture medium is removed, and fresh culture medium is added to the cells. The culture dishes are then  
25   returned to the rocking environment. Cells will also remain in suspension when cultured in suspension culture dishes (Nunc) without rocking, or when cultured in the absence of serum, which provides attachment factors. All cells must be cultured at 37°C, in a humidified,  
30   controlled gas atmosphere (either 5% CO<sub>2</sub>, 5% O<sub>2</sub> and 90% N<sub>2</sub> or 5% CO<sub>2</sub> in air).

          Following culture in suspension for up to 11 days, embryoid bodies are dispensed by mechanical or chemical means and can be allowed to reattach to tissue culture  
35   plates treated with gelatin or matrix, in ES medium.



Displaced, plated embryoid bodies will form flattened monolayers and can be maintained by replacing medium every 2 days.

Analysis Of Embryoid Bodies And Differentiated Cells

5 We used immunofluorescent antibody staining up to 7 days after plating to confirm the existence of cells of the neural phenotype. Cells were fixed in 30% methanol/10% acetic acid before incubation with antibodies. Antibodies that were used are as follows:  
10 rabbit anti-bovine GFAP (DAKO), anti-Forse-1 (Developmental Studies Hybridoma Bank), anti-bovine MAP-2 (Roche), anti-human NCAM/CD56 (DAKO) and anti-O1 (provided by S.-C. Zhang, University of Wisconsin). All primary antibodies are mouse monoclonals except anti-  
15 GFAP. Secondary antibodies, FITC-conjugated goat anti-mouse IgG and biotin-conjugated goat anti-rabbit, as well as AMCA-conjugated streptavidin were purchased from Jackson ImmunoResearch.

The Forse-1 antibody recognizes phosphacan, a brain-specific chondroitin sulfate proteoglycan that binds  
20 neural cell adhesion molecules in the embryonic CNS of both humans and rodents. K. Allendorfer *et al.*, 6 Mol. And Cell. Neuro. 381-395 (1995); S. Tole *et al.*, 15 J. Neuro. 957-969 (1995). The O1 antibody identifies pro-  
25 oligodendrocytes present from day 3 in embryonic mouse brain cultures. M. Schachner *et al.*, 83 Dev. Biol. 328-338 (1981); I. Sommer, *et al.*, 83 Dev. Biol. 311-327 (1981).

Within three days of plating, the neural precursors,  
30 stained by Forse-1 and O1, were observed. The Forse-1 antibody stained numerous rounded cells, whereas very sparse, flattened cells with extensive projections were stained with the anti-O1 antibody.

Neurons and glial were detected, 3 days and later  
35 after plating, by positive staining of neural cell

adhesion molecule (NCAM)/CD56 (Figure 3), microtubule-associated protein-2 (MAP-2) (Figure 3),  $\beta$ III-tubulin and glial fibrillary acidic protein (GFAP). NCAM is a cell adhesion molecule thought to be important in cell-cell interactions within the neuroepithelium. B. Cunningham, et al., 236 Science 799-806 (1987); J. Ritz, et al., 42 Adv. Immuno. 181-211 (1988). MAP-2 plays an important role in brain microtubule assembly.  $\beta$ III-tubulin is a neuron-specific marker, and glial fibrillary acidic protein (GFAP) is an astrocyte marker.

#### Differentiation Into Lineages

Embryoid bodies can be differentiated into a variety of desired lineages. For example the embryoid bodies could be used to derive hematopoietic cells using techniques analogous to those used for mouse in M. Wiles et al. 111 Development 259-267 (1991). In this regard one could plate the embryoid bodies in serum-containing medium in the presence of 2 i.u./ml erythropoietin or IL-3.

If cardiac lineages are desired one could use techniques analogous to T. Doetschman et al., 87 J. Embry. Exper. Morph. 27-45 (1985). One could plate the bodies in serum-containing medium with no additives.

To develop neural lineages one could plate the embryoid bodies in the presence of 20 ng/ml fibroblast growth factor plus 20 ng/ml epidermal growth factor. This is analogous to techniques described in B. Reynolds et al., 255 Science 5052 (1992).

The present invention thus provides an effective method for making primate embryoid bodies from primate ES cells. While the above work was focused on rhesus and human embryonic stem cells (and neural cells derived therefrom via these embryoid bodies), the techniques described herein should work broadly for primate embryonic stem cells and other cell types. Further,

while specific techniques for clump removal have been discussed, the invention is not limited to those alone. Rather, other techniques for removing the cells in clumps from the substrate should work.

5           Thus, the invention is not limited to the specific embodiments described herein. Rather, the claim should be looked to in order to judge the full scope of the invention.

#### Industrial Applicability

10           The present invention provides a supply of human and other primate embryoid bodies suitable for research, medical purposes, and differentiation into lineages.

CLAIMS

We claim:

1. A method for producing primate embryoid bodies from colonies of primate embryonic stem cells that are adhering to a substrate, the method comprising:  
removing the adhering colonies of the embryonic stem  
5 cells from the substrate in clumps; and  
then incubating the clumps in a container under conditions in which the clumps are essentially inhibited from attaching to the container and coalesce into embryoid bodies.
2. The method of claim 1, wherein the removal step is conducted in the presence of an enzyme that promotes disassociation of the clumps as clumps from the substrate.
3. The method of claim 2, wherein the enzyme is dispase.
4. The method of claim 1, wherein the removal step is conducted in the presence of a chelating agent.
5. The method of claim 1, wherein the removal step comprises mechanically scraping the clumps from the substrate.
6. The method of claim 1, wherein the removal step is conducted in the presence of trypsin, calcium and magnesium.
7. The method of claim 1, wherein the incubation step comprises agitating the container.
8. The method of claim 1, wherein the incubation step is conducted in a container made of plastic.
9. The method of claim 1, wherein the incubation step is conducted in the presence of a serum-free medium.
10. The method of claim 1, wherein the primate embryonic stem cells are human embryonic stem cells and the primate embryoid bodies are human embryoid bodies.

11. A primate embryoid body derived from the method of claim 1.

12. The primate embryoid body of claim 11, wherein the embryoid body is a human embryoid body.

13

13. A differentiated primate cell derived from the embryoid body of claim 11.

14. The differentiated primate cell of claim 13, wherein the cell is a human cell.

15. The differentiated primate cell of claim 14, wherein the cell is a human neural cell.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/62899 A3**

(51) International Patent Classification<sup>7</sup>: C12N 5/06, 5/00

(21) International Application Number: PCT/US01/05252

(22) International Filing Date: 20 February 2001 (20.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/510,444 21 February 2000 (21.02.2000) US

(71) Applicant: WISCONSIN ALUMNI RESEARCH  
FOUNDATION [US/US]: 614 Walnut Street, P O Box  
7365, Madison, WI 53707-7365 (US).

(72) Inventors: THOMSON, James, A.: 1807 Regent Street,  
Madison, WI 53705 (US). MARSHALL, Vivienne,  
S.: 4326 Buckeye Road, Madison, WI 53716 (US).  
SWIERGIEL, Jennifer, J.: 10219 Hawks Point Trail,  
Roscoe, IL 61073 (US).

(74) Agent: SEAY, Nicholas, J.: Quarles & Brady LLP, P O  
Box 2113, Madison, WI 53701-2113 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

(88) Date of publication of the international search report:  
21 February 2002

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHOD OF MAKING EMBRYOID BODIES FROM PRIMATE EMBRYONIC STEM CELLS

(57) Abstract: Primate embryoid bodies are formed from primate ES cells. The ES cells form clumps. One then removes the clumps, as clumps, and permits incubation under non-adherent conditions. The development of embryoid bodies from primate ES cells is dependent on maintaining the aggregation of cells, as individualized cells will rapidly die.

WO 01/62899 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/05252

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N5/06 C12N5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ITSKOVITZ-ELDOR J ET AL: "DIFFERENTIATION OF HUMAN EMBRYONIC STEM CELLS INTO EMBRYOID BODIES COMPRISING THE THREE EMBRYONIC GERM LAYERS" MOLECULAR MEDICINE, BLACKWELL SCIENCE, CAMBRIDGE, MA, US, vol. 6, February 2000 (2000-02), pages 88-95, XP000942869 ISSN: 1076-1551 the whole document --- -/--	1,2,6, 8-15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

18 September 2001

Date of mailing of the international search report

25/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/05252

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THOMSON J A ET AL: "EMBRYONIC STEM CELL LINES DERIVED FROM HUMAN BLASTOCYSTS" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE,, US, vol. 282, 6 November 1998 (1998-11-06), pages 1145-1147, XP002121340 ISSN: 0036-8075 cited in the application the whole document ---	1-4,8, 10-15
A	WO 99 27076 A (LORING JEANNE F ;ARC GENOMIC RESEARCH (US)) 3 June 1999 (1999-06-03) abstract page 16, line 5-20; examples 3,4 ---	1,6,13
A	US 5 843 780 A (THOMSON JAMES A) 1 December 1998 (1998-12-01) cited in the application column 7, line 38-53 column 8, line 28-67 ---	1,2,6,8, 11,13
A	DE 197 56 864 C (BRUESTLE OLIVER DR) 29 April 1999 (1999-04-29) page 2, line 3-5 page 4, line 11-17; examples 1-3 ---	1,6,8, 10-15
A	US 5 670 372 A (HOGAN BRIGID L M) 23 September 1997 (1997-09-23) column 2, line 6-19 column 5, line 55 -column 6, line 49 ---	1,2,6,8
A	TOLE SHUBHA ET AL: "FORSE-1: A positionally regulated epitope in the developing rat central nervous system." JOURNAL OF NEUROSCIENCE, vol. 15, no. 2, 1995, pages 957-969, XP001019798 ISSN: 0270-6474 cited in the application abstract ---	1,10,15
A	SOMMER I ET AL: "MONO CLONAL ANTIBODIES 0-1 TO 0-4 TO OLIGODENDROCYTE CELL SURFACES AN IMMUNO CYTOLOGICAL STUDY IN THE CENTRAL NERVOUS SYSTEM" DEVELOPMENTAL BIOLOGY, vol. 83, no. 2, 1981, pages 311-327, XP001020302 ISSN: 0012-1606 cited in the application abstract -----	1,10,15

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/05252

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9927076	A	03-06-1999	AU	1704899 A	15-06-1999
			CN	1284993 T	21-02-2001
			EP	1060244 A1	20-12-2000
			WO	9927076 A1	03-06-1999
US 5843780	A	01-12-1998	US	6200806 B1	13-03-2001
			AU	4758496 A	07-08-1996
			CA	2190528 A1	25-07-1996
			EP	0770125 A1	02-05-1997
			WO	9622362 A1	25-07-1996
DE 19756864	C	29-04-1999	DE	19756864 C1	29-04-1999
			AU	2510699 A	12-07-1999
			WO	9932606 A2	01-07-1999
			EP	1040185 A2	04-10-2000
US 5670372	A	23-09-1997	US	5453357 A	26-09-1995
			US	5690926 A	25-11-1997